

GENERAL INSTRUCTIONS TO THE USERS

The Blulux Dehydrated Culture medias are highly hygroscopic and get deteriorated easily unless stored in a cool & dry place away from bright light. So care must be taken in the storage of dehydrated media.

Following instructions should be followed while using dehydrated culture media.

- ★ Read carefully the instruction given on the label.
- ★ Note the batch no. and date before opening the container.
- ★ Before use confirm that the media is not deteriorated physically.
- ★ Since the media is hygroscopic in nature, please ensure that the container is tightly closed and stored in cool and dry place after use.

DISSOLVING THE MEDIA

For dissolving use clean undamaged glassware two-three times larger in volume than the final volume of media to be prepared. Water which meets the U.S.P./I.P specifications should only be used.

Place accurately weighed amount of the medium in a clean dry flask. Add part of water and swirl until dissolved then add the remaining water through the sides of the flask to make up the required volume. For complete dissolution heat the media taking care to avoid overheating or scorching.

ADJUSTMENT OF pH

After preparation the pH of the media should be in concordance with value mentioned on the label at 25 °C. If required, the pH should be adjusted to the specified value by adding 1N or 0.1N HCl or NaOH solution.

STERILIZATION

Sterilization of the media is usually carried out at 121 °C for 15minutes using an autoclave. Autoclave efficiency should be checked from time to time.

POURING OF STERILIZED MEDIA

After sterilization Agar media should be poured into petridishes at 45-50 °C. The medium should be mixed well avoiding bubble formation. Agar surface should be dried at 30-40 °C in the incubator before inoculation to avoid microbial swarming.

STORAGE OF PREPARED MEDIA

Unless the medium is not used in the same day it is prepared, then it should be kept in moisture proof containers.

Agar containing media should not be stored at higher temperatures. Agar plates should be stored at 2-8 °C in sealed moisture proof containers. Stability of the prepared culture media is limited and varies considerably. Never store media below 0 °C as it destroys its gel strength.

DISPOSAL OF MEDIA

All specimens and cultures should be carefully handled and must not disposed without autoclaving. Cultures in vessels should be autoclaved for approximately 30 minutes at 121 °C before disposal.

STORAGE OF CULTURE MEDIA PRODUCTS

For obtaining desired results **Blulux Culture Media** products should be stored in specified storing conditions. It is recommended to use the products in the order of Batch number/Mfg. date.

Storage temperature and shelf-life of the *Blulux* Culture Media products are as follows.

DEHYDRATED MEDIA

Blulux Dehydrated Culture Media if stored under specified conditions will have shelf-life of 2 to 3 years. Storage temperature for dehydrated media are preferably between 8 and 20°C.

CERTAIN PRECAUTIONS WHILE USING BLULUX DEHYDRATED MEDIA

FAULTS	REASONS
Drift in pH	Overheating, incomplete mixing, prolonged sterilization, use of alkaline glass, impure water, hydrolysis of ingredients, prolonged storage at high temperature.
Incomplete Solubility	Inadequate heating of agar media incomplete mixing.
Darkening	Overheating of the medium, excess amount of dehydrated powder, improper mixing.
Soft gel	Agar not in solution, improper reconstitution of dehydrated medium, acid hydrolysis of agar, failure to compensate for dilution of agar by the inoculum.
Loss of growth promoting of tiating properties	Repeated remelting, excessive heating, incomplete mixing, failure to compensate for differendilution of ingredients, disturbance in the formula by inoculum carriers, etc.
Abnormal colour of Medium	Deteriorated Dehydrated Medium improperly washed glassware impure water.
Contamination	Improper/Insufficieent sterilization. Poor technique in adding enrichments and pouring plates.

Please Contact us Or call us for custom synthesis & technical information / inquiries on any of **Blulux Culture Medias** Products or for the availability of specific **Blulux Dehydrate Culture Media** not currently in the price list.

Please E-mail us at blulux@rediffmail.com & bluluxlab@gmail.com

**We Offer
Special Prices
For Exports & Bulk Requirments**

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1A001	A.1 Medium (Broth) For determining the presence of faecal coliforms in water samples and foods by MPN technique.	31.50 gms/ltr.	703	2827
1A002	AC Agar For cultivation of a wide variety of microorganisms particularly for sterility testing.	35.20 gms/ltr.		1881
1A003	AC Broth for cultivation of common aerobes and sterility testing of solutions and biological products without mercurial preservatives.	34.20 gms/ltr.		1773
1A004	*Acetate Agar for the isolation and cultivation of <i>Leuconostoc</i> and <i>Pediococcus species</i> .	61.90 gms/ltr.		2054
1A005	Acid Broth For the isolation of bacteria from canned foods.	27.50 gms/ltr.		1728
1A006	Alkaline Peptone Water for enrichment of <i>Vibrio species</i> .	20.00 gms/ltr.	360	1401
1A007	Alkaline Peptone Water recommended by ISO Committee for enrichment of <i>Vibrio species</i> .	50.00 gms/ltr.		1401
1A008	Alkaline Peptone Water recommended by BIS for enrichment of <i>Vibrio species</i> .	15.00 gms/ltr.	320	1465
1A009	Alternative Thioglycollate Medium (Thioglycollate Broth, Alternative) for sterility testing of turbid or viscous biological products.	29.00 gms/ltr.	183	699
1A010	Amies Transport Medium w/ Charcoal for transportation and preservation of microbiological specimens.	20.00 gms/ltr.	423	1688
1A011	Anaerobic Agar a general purpose medium for the cultivation of anaerobic bacteria, especially, <i>Clostridium species</i> .	58.00 gms/ltr.		1886
1A012	Anaerobic Agar (Brewer) for the isolation and sensitivity testing of anaerobic and microaerophilic organisms and study of colonial morphology.	53.00 gms/ltr.		1886
1A013	* Anaerobic CNA Agar Base for the selective isolation of anaerobic <i>Streptococci</i> .	44.14 gms/ltr.	3734	
1A014	Anaerobic Egg Agar Base for detection of <i>Clostridium perfringens</i> in food samples.	55.00 gms/ltr.		1998
1A015	Andrade Peptone Water recommended by BIS as a basal medium, which with carbohydrate addition is used to study fermentation reactions.	15.10 gms/ltr.	351	1617
1A016	Andrade Peptone Water a basal medium, which with carbohydrate addition is used to study fermentation reactions.	15.10 gms/ltr.	336	1515
1A017	Antibiotic Assay Medium B for microbiological assay of colistin sulphate and polymyxin B sulphate using <i>Bordetella bronchiseptica</i> ATCC 4617 in accordance with B.P.	55.00 gms/ltr.		2019
1A018	Antibiotic Assay Medium C for microbiological assay of antibiotics using <i>Staphylococcus aureus</i> NCTC 6571.	20.00 gms/ltr.		2013
1A019	Antibiotic Assay Medium E for microbiological assay of Neomycin sulphate and Framycetin sulphate using <i>Bacillus subtilis</i> ATCC 6633 and <i>Bacillus pumilus</i> NCTC 8241 in accordance with B.P.	28.70 gms/ltr.		2019

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1A020	Antibiotic Assay Medium No.1 (Seed Agar) For microbiological assay of b-lactam and other antibiotics in accordance with USP/IP.	30.50 gms/ltr.	442	1769
1A021	Antibiotic Assay Medium No.2 (Base Agar) for microbiological assay of antibiotics in accordance with USP/IP.	25.50 gms/ltr.		2025
1A022	Antibiotic Assay Medium No.3 for microbiological assay of antibiotics in accordance with USP/IP.	17.50 gms/ltr.	378	1698
1A023	Antibiotic Assay Medium No.4 (Yeast Beef Agar) for detection of Penicillin-G in milk samples using <i>Bacillus stearotherophilus</i> .	26.50 gms/ltr.		2039
1A024	Antibiotic Assay Medium No.5 (Streptomycin Assay Agar w/Yeast Extract) microbiological assay of Streptomycin using <i>Bacillus subtilis</i> , in accordance with USP/IP.	25.50 gms/ltr.		2019
1A034	Antibiotic Assay Medium No. 8 (Base Agar w/low pH) for microbiological assay of Mitomycin, Plicamycin & Vancomycin, in accordance with USP/IP.	25.50 gms/ltr.		1926
1A025	Antibiotic Assay Medium No. 10 (Polymyxin Seed Agar) as a seed layer medium for assaying the products containing Polymxin-B, also for assaying Carbenicillin, Colistin and Colistimethate sodium.	52.00 gms/ltr.		2019
1A026	Antibiotic Assay Medium No. 11 (Neomycin, Erythromycin Assay Agar) (Erythromycin Seed Agar) for microbiological assay of antibiotics in accordance with USP.	30.50 gms/ltr.	442	1769
1A027	Antibiotic Assay Medium No. 19 for microbiological assay of Amphotericin B, Netamycin & Nystatin using <i>Saccharomyces cerevisiae</i> ATCC 9763 & 2601, in accordance with USP/IP.	60.00 gms/ltr.		2095
1S019	Antibiotic Assay Medium No. 36 (Tryptone Soya Agar) (see Soyabean Casein Digest Agar)	40.00 gms/ltr.	384	1539
1S020	Antibiotic Assay Medium NO. 37 (Tryptone Soya Broth) (see Soyabean Casein Digest Medium)	30.00 gms/ltr.	183	699
1A028	Antibiotic Assay Medium No. 38 for microbiological assay of Ticarcil/in using <i>Pseudomonas aeruginosa</i> ATCC 29336.	45.40 gms/ltr.		2019
1A029	Antifungal Assay Agar for assaying antifungal activity of pharmaceutical products and other materials by the cylinder plate or disc method.	75.70 gms/ltr.		1938
1A030	AK Agar No. 2 (Sporulating Agar) (Arret & Klrshbaum Medium) for production of spores of <i>Bacillus subtilis</i> ATCC 6633 which are used as inoculum in detection of Penicillin & other antibiotics.	30.80 gms/ltr.		1936
1A031	Ashby's Glucose Agar for cultivation of <i>Azotobacter</i> species that can use glucose/mannitol and Atmospheric nitrogen as source of carbon and nitrogen respectively.	40.70 gms/ltr.		1998
1A032	Ashby's Mannitol Agar for isolation of <i>Azotobacter</i> species from soil	40.70 gms/ltr.		1998
1A035	Ayers and Johnson Agar (Stock Culture Agar) for maintenance of cultures of <i>Streptococci</i> and other microorganisms.	50.00 gms/ltr.		1869
1A033	Azide Dextrose Broth w/ BCP recommended by ISO committee for cultivation of faecal <i>Streptococci</i> .	34.70 gms/ltr.		1974
1B001	B.T.B. Lactose Agar For isolation of pathogenic <i>Staphylococci</i>	33.17 gms/ltr.		1938

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1B011	*B12 Assay Medium (*Vitamin B12 Assay Medium) for microbiological assay of vitamin B ₁₂ using <i>Lactobacillus leichmannii</i> ATCC 7830 as the test organism.	85.00 gms/ltr.	1594	
1B002	*B12 Culture Agar (E. Coli Maintenance Medium) (E. Coli Mutant Culture Agar) for propagation, cultivation and maintenance of <i>Escherichia coli</i> mutant used in microbiological assay of Vitamin B ₁₂ .	32.80 gms/ltr.	879	
1B003	*B12 Culture Agar (L. Leichmannii Maintance Medium) for propagation cultivation and maintenance of <i>Lactobacillus leichmannii</i> ATCC 7830.	42.10 gms/ltr.	653	
1B004	*B12 Inoculum Broth for preparing the inoculum of <i>Lactobacillus leichmannii</i> ATCC 7830 for the microbiological assay of Vitamin B ₁₂ .	32.10 gms/ltr.	653	
1B005	BPL Agar For isolation and identification of <i>Salmonellae</i> except <i>Salmonella typhi</i> in faeces, urine, milk and other materials.	40.00 gms/ltr.		1869
1B006	Bacillus Cereus Agar Base a selective medium for isolation, detection and enumeration of <i>Bacillus cereus</i> .	41.00 gms/ltr.	489	1962
1B007	Baird Parker Agar Base for isolation & enumeration of coagulase positive <i>Staphylococci</i> from food & other materials.	63.00 gms/ltr.	516	2070
1B008	Baird Parker Agar Base recommended by BIS and enumeration of coagulase positive <i>Staphylococci</i> from food & other materials.	65.00 gms/ltr.	516	2070
1B009	Baird Parker Agar Base for isolation and enumeration of coagulase positive <i>Staphylococci</i> from food and other materials in accordance with I.P.	63.00 gms/ltr.	516	2070
1B010	Baird Parker Agar Base w/ Sulpha recommended for isolation and enumeration of coagulase positive <i>Staphylococci</i> from food and other materials.	63.00 gms/ltr.		2367
1A021	Base Agar (see Antibiotic Assay Medium No.2)	25.50 gms/ltr.		2025
1A034	Base Agar w/low pH (see Antibiotic Assay Medium No. 8)	25.50 gms/ltr.		1945
1B013	Beef Extract Paste			523
1B014	Beef Extract Powder refined for use in microbial culture media			1225
1B015	Beef Extract Powder, Certified for the maximum recovery and growth of a wide variety of microorganisms.			2300
1B016	Beef Extract Powder Type 1 used in media for routine cultivaation & diagnostic purposes.			806
1B017	Bi. G.G.Y. Agar (Nickerson Medium) for detection, selective isolation, differentiation and presumptive identification of <i>Candida albicans</i> and <i>Candida tropicalis</i> .	45.00 gms/ltr.		1917
1B018	Bile Esculin Agar for differential isolation and presumptive identification of group D <i>Streptococci</i> in food & pharmaceutical products.	64.50 gms/ltr.	1077	3497
1B019	Bile Esculin Agar recommended by ISO Committee for isolation and identification of <i>Yersinia enterocolitica</i> .	64.50 gms/ltr.		3497

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1B020	Bile Esculin Agar Base for differential isolation and presumptive identification of group D <i>Streptococci</i> in Food & pharmaceutical products.	63.50 gms/ltr.		2268
1B021	Bile Salts for Bacteriology.		1260	3780
1B022	Bile Salts, Certified a specially manufactured extract of bile salts, prepared from fresh ox bile, is recommended as a selectively inhibitory agent in microbiological culture media.			4047
1B023	Bile Salt Agar for isolation and enumeration of bile tolerant enteric bacilli.	43.00 gms/ltr.	459	2037
1B024	Bile Salt Agar recommended by BIS for isolation and enumeration of bile tolerant enteric bacilli.	40.00 gms/ltr.	474	2037
1B025	Bile Salt Brilliant Green Starch Agar for selective isolation & identification of <i>Aeromonas hydrophila</i> from food & Environmental specimens.	45.00 gms/ltr.		2037
1B026	Bile Salts Mixture Equivalent to Bile Salt No.3		2240	
1B027	Bile Salts Mixture, Certified recommended for use in microbiological culture media for selective isolation and cultivation of bile tolerant enteric bacteria.			4394
1B028	Bio Peptone mixture of Casein and Meat Peptones employed in media used for cultivation of microorganisms.			1324
1B029	Bio Peptone, Certified provides a broad spectrum of peptides & amino acids which supports better microbiological growth characteristics to a large variety of organisms.			2324
1B030	Bismuth Sulphite Agar for selective isolation and identification of <i>Salmonellae</i> in accordance with I.P.	63.40 gms/ltr.	424	1680
1B031	Bismuth Sulphite Agar for selective isolation of <i>Salmonellae</i> from faeces, urine, sewage and other materials. in accordance with U.S.P.	52.32 gms/ltr.	433	1725
1B032	Bismuth Sulphite Agar for selective isolation of <i>Salmonellae</i> from faeces, urine, sewage and other materials.	52.33 gms/ltr.	348	1395
1B033	Blood Agar Base (Infusion Agar) for isolation and cultivation of many fastidious pathogenic microorganisms after addition of blood.	40.00 gms/ltr.	348	1395
1B034	Blood Agar Base No.2 After addition of blood, medium permits maximum recovery of <i>Streptococci</i> , <i>Pneumococci</i> and other fastidious pathogenic microorganisms without interfering with their haemolytic reactions.	42.50 gms/ltr.	348	1395
1B035	Blood Agar Base No.2 w/1.2% Agar Especially devised to permit the maximum recovery of fastidious pathogenic Microorganisms without interfering with their haemolytic reactions.	39.50 gms/ltr.		1566
1B036	Blood Free Campylobacter Selectivity Agar Base recommended by ISO Committee for selective isolation & differentiation of <i>Campylobacter species</i> .	45.50 gms/ltr.		2057

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1B037	Bordet Gengou Agar Base for detection and isolation of <i>Bordetella pertussis</i> and <i>Bordetella parapertussis</i> .	40.00 gms/ltr.		1944
1B038	Bordet Gengou Agar Base w/1.6% Agar for detection and isolation of <i>Bordetella pertussis</i> and <i>Bordetella parapertussis</i> .	36.00 gms/ltr.		1944
1B039	Brain Heart Infusion Powder highly nutritious ingredient, used for cultivation of fastidious microorganisms.			1477
1B040	Brewer Thioglycollate Medium for testing the sterility of biological products and for the isolation of aerobic and anaerobic organisms	40.50 gms/ltr.	264	1185
1B041	Brewer Thioglycollate Medium, Modified (Linden Thioglycollate Medium) for testing sterility of biological products and for the isolation of aerobic and anaerobic organisms.	38.50 gms/ltr.	264	1185
1B042	Brewer Thioglycollate Medium, Modified for testing sterility of biological products and for isolation of aerobic and anaerobic organisms.	20.00 gms/ltr.	264	1185
1B043	Brilliant Green Agar, Modified for selective isolation of <i>Salmonellae</i> other than <i>Salmonella typhi</i> from faeces, foods, dairy products etc.	58.00 gms/ltr.	348	1395
1B044	Brilliant Green Agar, Modified for selective isolation of <i>Salmonellae</i> other than <i>Salmonella typhi</i> from faeces, foods, dairy products etc. in accordance with U.S.P.	58.09 gms/ltr.	338	1521
1B045	Brilliant Green Agar w/ 1.2% Agar for selective isolation of <i>Salmonellae</i> other than <i>Salmonella typhi</i> from faeces, foods, Dairy products etc.	50.00 gms/ltr.	352	1413
1B046	Brilliant Green Agar Base w/ Phosphates recommended by ISO committee for selective isolation of <i>salmonellae</i> while inhibiting <i>Escherichia coli</i> , <i>Proteus</i> and <i>Pseudomonas</i> species.	52.00 gms/ltr.		1787
1B047	Brilliant Green Agar w/Phosphates recommended by ISO committee for isolation of <i>Salmonellae</i> while inhibiting <i>Escherichia coli</i> <i>Proteus</i> and <i>Pseudomonas species</i> .	54.69 gms/ltr.		1884
1B048	Brilliant Green Agar Base w/Phosphates recommended by BIS for selective isolation of <i>Salmonella</i> while inhibiting <i>Escherichia coli</i> , <i>Proteus</i> and	52.00 gms/ltr.	417	1884
1B049	Brilliant Green Bile Broth 2% recommended by Iso Committee for detection & confirmation of coliform bacteria in water, waste water, foods, milk and dairy products.	40.00 gms/ltr.	348	1395
1B050	Bromo Cresol Purple Borth for studing fermentation of carohydrates by pure cultures	18.00 gms/ltr.		1938
1B051	Brucella Agar Base for selective isolation and cultivation of <i>Brucella</i> or <i>Campylobacter</i> species from Clinical & nonclinical specimens.	43.10 gms/ltr.	433	1725
1B052	Brucella Agar Base, Modified for cultivation of <i>Campylobacter</i> species.	44.10 gms/ltr.		1725
1B053	Brucella Agar w/ Hemin and Vitamin K for cultivation of <i>Brucella</i> species, for isolation & subculture of anaerobes with Addition of blood.	43.00 gms/ltr.		2102

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1B054	Brucella Broth Base for enrichment and cultivation of <i>Brucella</i> or <i>Campylobacter</i> species from clinical and nonclinical specimens.	28.10 gms/ltr.	433	1725
1B055	Brucella Selective Medium Base for isolation and identification of <i>Brucella</i> species.	43.50 gms/ltr.		1881
1B056	Buffered Charcoal Yeast Extract Agar Base for selective isolation and cultivation of <i>Legionella</i> species from clinical and other materials.	40.00 gms/ltr.	2228	
1B060	Buffered Peptone Water as a pre-enrichment medium for the isolation of <i>Salmonella species</i> from foods products.	20.00 gms/ltr.	266	1302
1B057	Bushnell Haas Agar for examination of fuels for microbial contamination and for studying hydrocarbon Deterioration by microorganisms.	23.27 gms/ltr.		1869
1B058	Bushnell Haas Broth for examining fuels for microbial contamination and for studying hydrocarbon Deterioration by microorganisms.	3.27 gms/ltr.	392	1765
1C001	C.L.E.D. Agar w/ Andrade Indicator for isolation, and differentiation of urinary pathogens on the basis of lactose fermentation.	36.25 gms/ltr.	381	1526
1C002	C.L.E.D. Agar Base w/o Indicator for isolation, enumeration and presumptive identification of bacterial flora in urinary tract.	36.10 gms/ltr.		1617
1C003	C.L.E.D. Agar w/Bromo Thymol Blue for isolation and differentiation of urinary pathogens on the basis of lactose fermentation.	36.15 gms/ltr.	366	1580
1C004	Cary-Blair Medium Base (Transport Medium w/o Charcoal) for collection and shipment of clinical specimens.	12.60 gms/ltr.	492	2093
1C005	Casein Acid Hydrolysate Vitamin Free			4433
1C006	Cetrimide Agar Base for selective isolation of <i>Pseudomonas aeruginosa</i> from clinical specimen.	46.70 gms/ltr.	360	1481
1C007	Cetrimide Agar Base w/o Glycerine selective isolation of <i>Pseudomonas aeruginosa</i> in accordance with I.P.	45.30 gms/ltr.	321	1467
1C008	Cetrimide Broth for selective cultivation of <i>Pseudomonas aeruginosa</i>	25.30 gms/ltr.	360	1481
1C009	Cetrimide Broth Base recommended by ISO Committee for cultivation of <i>Pseudomonas aeruginosa</i> from water samples using membrane filter technique.	31.90 gms/ltr.		1796
1C010	Chapman Stone Agar selective isolation of <i>Staphylococci</i> causing food poisoning,	202.50 gms/ltr.		1796
1C011	Charcoal Agar Base for cultivation of <i>Bordetella pertussis</i> , for vaccine Production and also for stock culture maintenance.	62.50 gms/ltr.		1938
1C012	Chlamydospore Agar for differentiation of <i>Candida albicans</i> from other <i>Candida</i> species on the basis of chlamydospore formation.	37.10 gms/ltr.	666	2997
1C013	Chocolate Agar Base for isolation of <i>Neisseria gonorrhoeae</i> from chronic & acute cases of gonococcal infections.	45.00 gms/ltr.		1918

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. MEDIUM	100 GM	500 GM
1C030	Chocolate Agar an enriched medium for the isolation & cultivation of fastidious organisms as neisseria & haemophilus species. Haemoglobin & Bio-X. Enrichment replace lysed heated Blood & give to the culture media enough nutrients agents to permit luxuriant growth of fastidious microorganisms.	36.00 gms/ltr.		3033
1C014	Citrate Agar for cultivation of iron bacteria from soil samples	27.20 gms/ltr.	330	1485
1C015	Clausen Medium for sterility testing of Nordic Pharmacopoeia Board.	40.00 gms/ltr.		1875
1C016	Clostridium Difficile Agar Base for selective isolation of <i>Clostridium difficile</i> from faeces.	69.10 gms/ltr.		1884
1C017	Columbia Blood Agar Base an efficient base for preparation of blood agar, chocolate agar and for various selective and identification media.	44.00 gms/ltr.	417	1689
1C018	Columbia Blood Agar Base w/1% Agar a basal medium used with or without blood for isolation & cultivation of fastidious bacteria.	39.00 gms/ltr.		1734
1C019	Columbia Broth Base cultivation of fastidious organisms from clinical specimens	35.00 gms/ltr.		1548
1C020	Columbia C.N.A. Agar for selective isolation of pathogenic gram-positive cocci from clinical and nonclinical specimens.	44.00 gms/ltr.		4260
1C021	Cooke Rose Bengal Agar Base for selective isolation and Cultivation of fungi	36.54 gms/ltr.		1938
1C022	Cooked Meat Medium (R.C. Medium) for cultivation of aerobes and anaerobes, especially pathogenic <i>Clostridia</i> and also for the maintenance of stock cultures.	125.00 gms/ltr.	433	1770
1C029	Corn Meal Agar for production of chlamydo spores by <i>Candida albicans</i> & the maintenance of fungal stock cultures.	17.00 gms/ltr.	363	1688
1C023	Crystal Violet Lactose Agar for differentiation of pure cultures of pathogenic and nonpathogenic <i>Staphylococci</i> .	33.00 gms/ltr.		1803
1C031	Culture Medium For RWC (Disinfectant Test Broth) for determination of phenol coefficients of disinfectants using <i>Salmonella typhi</i> as a test organism, in accordance with ISI specification.	50.00 gms/ltr.	456	1828
1C024	Czapek Dox Agar a semisynthetic medium for general cultivation of fungi.	49.00 gms/ltr.	330	1513
1C025	Czapek Dox Agar Modified for cultivation and maintenance of fungi.	45.36 gms/ltr.		1724
1C026	Czapek Dox Broth a semisynthetic medium used, for cultivation of fungi.	35.00 gms/ltr.	311	1270
1C027	Czapek Malt Agar for isolation, detection and cultivation of saprophytic fungi.	94.00 gms/ltr.	330	1513
1C028	Czapek Yeast Extract Agar for the cultivation and maintenance of <i>Aspergillus niger</i> .	61.00 gms/ltr.		1965

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. MEDIUM	100 GM	500 GM
1D003	D.C.L.S. Agar a selective medium used to detect & isolate <i>Salmonella</i> & <i>Shigella species</i> . Also Useful for isolation of <i>Virbio cholerae</i> .	49.50 gms/ltr.	330	1472
1D004	D.C.L.S. Agar, Hajna for the isolation of gram-negative enteric bacilli.	73.52 gms/ltr.	330	1472
1D005	Decarboxylase Broth Base, Moeller with the addition of appropriate L-amino acid, it is used to differentiate bacteria on the Basis of their ability to decarboxylate the amino acid.	10.52 gms/ltr.	1013	4077
1D006	Decarboxylase Test Medium Base (Falkow) recommended by BIS for testing amino acid decarboxylase activity.	9.00 gms/ltr.	396	1593
1D007	Decarboxylase Test Medium Base (Falkow) for testing amino acid decarboxylase activity.	9.00 gms/ltr.	393	1593
1D008	Deoxycholate Citrate Agar selective medium for the isolation of enteric pathogens particularly <i>Salmonella</i> and <i>Shigella species</i> .	70.52 gms/ltr.	280	1270
1D009	Deoxycholate Citrate Agar for selective isolation of enteric pathogen in accordance with B.P.	69.52 gms/ltr.		1270
1D010	Deoxycholate Citrate Agar for selective isolation and identification of Salmonellae in accordance with I.P.	55.00 gms/ltr.	343	1425
1D011	Deoxycholate Citrate Agar, Modified selective medium for the isolation of <i>Salmonella</i> & <i>Shigella species</i> .	52.00 gms/ltr.		1918
1D012	Dextrose Agar Base, Emmons (Sabouraud Dextrose Agar, Modified Emmons) for selective cultivation of pathogenic fungi.	47.00 gms/ltr.	442	1828
1D013	Dextrose Starch Agar for propagation of pure cultures of <i>Neisseria gonorrhoeae</i> and other fastidious Organisms.	65.00 gms/ltr.		1688
1D014	Dextrose Tryptone Agar for detection & enumeration of mesophilic & Thermophilic aerobic organisms in foods.	30.00 gms/ltr.		1688
1D015	Dextrose Tryptone Broth, Modified for isolation and cultivation of aciduric and thermophilic sporeformers.	17.30 gms/ltr.		1668
1D016	*Dey-Engley Neutralizing Agar used in disinfectant testing, where neutralization of the chemical is important for determining its bactericidal activity.	54.00 gms/ltr.		2093
1D017	*Dey-Engley Neutralizing Broth for neutralizing and testing antiseptics and disinfectants.	39.00 gms/ltr.		1974
1D018	Dey-Engley Neutralizing Broth Base used along with Dey-Engley Neutralizing Broth Base for neutralizing and determining bactericidal activity of quaternary ammonium compounds.	17.50 gms/ltr.		1688
1D024	Diagnostic Thloglycollate Medium (Thioglycollate Medium w/o Indicator) for enrichment of blood cultures.	30.00 gms/ltr.		1356
1D019	Differential Reinforced Clostridial Broth Base for the cultivation of <i>Clostridia</i> from water.	29.00 gms/ltr.	480	2004
1D020	Differential Reinforced Clostridial Broth Base recommended by ISO Committee for cultivation of <i>Clostridia</i> from water.	29.00 gms/ltr.		2004
1C031	Disinfectant Test Broth see Culture Medium for RWC	50.00 gms/ltr.	456	1828

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1D022	Disinfectant Test Broth, AOAC recommended by AOAC for testing disinfectants.	20.00 gms/ltr.		1701
1D001	DNase Test Agar Base for detection of deoxyribonuclease activity of bacteria and fungi, and especially for identification of pathogenic Staphylococci.	42.00 gms/ltr.	2822	
1D002	DNase Test Agar w/ Toluidine Blue detection of deoxyribonuclease activity of microorganisms.	42.10 gms/ltr.	3441	
1D023	Dubos Broth Base for preparation of liquid medium for rapid cultivation of pure cultures of <i>Mycobacterium tuberculosis</i> and related microorganisms.	6.50 gms/ltr.	424	1701
1B002	E. Coli Maintenance Medium (E. Coli Mutant Culture Agar) (See *B12 Culture Agar)	32.80 gms/ltr.	879	
1E001	EC Broth for selective enumeration of faecal & nonfaecal coliforms in water, wastewater & shell fish.	37.00 gms/ltr.	411	1648
1E002	EC Broth recommended by ISO Committee for selective enumeration of presumptive <i>Escherichia coli</i> by MPN technique	37.00 gms/ltr.		1648
1E003	EE Broth, Mossel for selective enrichment of Enterobacteriaceae in the bacteriological examination of foods.	43.50 gms/ltr.	459	1828
1E004	EE Broth Modified for selective enrichment of Enterobacteriaceae in the bacteriological examination of foods.	45.00 gms/ltr.		1884
1E005	EE Broth, Mossel for selective enrichment of Enterobacteriaceae in the bacteriological examination of foods.	45.00 gms/ltr.		1884
1E006	EE Broth, Mossel recommended by ISO Committee for selective enrichment of Enterobacteriaceae in bacteriological examination of foods.	43.47 gms/ltr.		1944
1E007	EE Broth, Mossel (Enrichment Broth Medium E) for selective enrichment of Enterobacteriaceae in the bacteriological examination of foods in accordance with E.P.	46.01 gms/ltr.		1905
1E014	Egg Yolk Agar Base for isolation and identification of <i>Clostridia</i> and certain other anaerobes.	75.00 gms/ltr.		1803
1E026	EGG Yolk Emulsion recommended for use in various culture media.	100 ml/vial	635/5vl	
1E008	EMB Agar for differential isolation of gram-negative enteric bacilli from clinical and nonclinical specimens.	36.00 gms/ltr.	348	1400
1E009	EMB Agar Base a basal medium to which different carbohydrates and other for differentiation and study of various enteric bacteria.	27.50 gms/ltr.	410	1650
1E010	EMB Agar, Levine recommended by BIS for isolation, enumeration and differentiation of members of Enterobacteriaceae.	37.50 gms/ltr.	388	1572
1E011	EMB Agar, Levine for isolation, enumeration and differentiation of members of Enterobacteriaceae in accordance with I.P.	37.50 gms/ltr.	388	1572

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. MEDIUM	100 GM	500 GM
1E012	EMB Agar, Levine for isolation, enumeration and differentiation of members of Enterobacteriaceae.	37.50 gms/ltr.	388	1572
1E013	EMB Broth for differentiation of gram-negative enteric bacteria.	22.50 gms/ltr.	330	1422
1E015	Eijkman Lactose Broth for detection and differentiation of <i>Escherichia coli</i> from other coliforms on the basis of their ability to grow and liberate gas from lactose.	28.50 gms/ltr.		1623
1E016	Elliker Broth (Lactobacilli Broth) for cultivation of <i>Lactobacilli</i> and <i>Streptococci</i> of importance in dairy industry.	48.50 gms/ltr.		1707
1E017	Endo Agar for the confirmation of the presumptive test for members of the coliform group.	41.50 gms/ltr.	338	1398
1E018	Endo Agar Base for preparing Endo Agar to confirm presumptive test for lactose fermenting coliforms.	38.00 gms/ltr.		1436
1E019	Enriched Thioglycollate Broth for isolation, cultivation and identification of a wide variety of obligate anaerobic bacteria.	30.00 gms/ltr.		1720
1E020	Enterococcus Presumptive Broth for detection of <i>Enterococci</i> in water and other materials of sanitary importance.	15.40 gms/ltr.		1701
1E027	Enzymic Digest of Casein			1930
1E028	Enzymic Digest of Gelatin			1540
1A026	Erythromycin Seed Agar (Neomycin, Erythromycin Assay Agar) (see Antibiotic Assay Medium No.11)	30.50 gms/ltr.	442	1769
1E022	Esculin Agar for cultivation and differentiation of bacteria that can hydrolyze esculin and produce H ₂ S	41.50 gms/ltr.		2094
1E023	Eugonic Agar cultivation of fastidious microorganisms like <i>Haemophilus</i> , <i>Neisseria</i> , <i>Pasteurella</i> , <i>Brucella</i> and <i>Lactobacillus</i> species.	44.40 gms/ltr.		1801
1E024	Eugonic Broth cultivation of fastidious microorganisms like <i>Haemophilus</i> , <i>Neisseria</i> , <i>Pasteurella</i> , <i>Brucella</i> and <i>Lactobacillus</i> species.	29.40 gms/ltr.		1701
1F015	Fermentation Broth Base Supplemented with the appropriate carbohydrates as used to determine the fermentation reactions of microorganisms	16.00 gms/ltr.	537	2140
1F001	*Fluid Casein Digest Soya Lecithin Medium (Twin Pack) For sanitary examination of surfaces.	25.00 gms/ltr.	1000	3765
1F002	Fluid Lactose Medium as a pre enrichment medium for detection of coliform bacteria in water, dairy Products and foods.	13.00 gms/ltr.	237	984
1F003	Fluid Sabouraud Medium (Sabouraud Medium, Fluid) sterility test medium for moulds and lower bacteria in pharmaceutical preparations.	30.00 gms/ltr.	190	748
1F005	Fluid Selenite Cystine Broth recommended by ISO Committee as an enrichment medium for isolation of <i>Salmonellae</i> from faeces, urine or other pathological materials.	17.00		1860
1F004	Fluid Selenite Cystine Medium (Selenite Cystine Broth) enrichment medium for isolation of <i>Salmonellae</i> in foods, dairy products and materials of sanitary importance and clinical specimens.	23.00 gms/ltr.	498	1999

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1F006	Fluid Tetrathionate Medium w/o Iodine and BG (Tetrathionate Broth Base w/o Iodine and BG) selective enrichment medium for isolation of <i>Salmonellae</i> from foods and other pathological materials.	46.00 gms/ltr.	248	1080
1F007	Fluid Thioglycollate Medium (Thioglycollate Medium Fluid) for sterility testing of biologicals and for cultivation of aerobes, anaerobes and microaerophiles in accordance with USP.	29.75 gms/ltr.	181	699
1F008	Fungal Agar (Mycological Agar) for cultivation and maintenance of fungi	35.00 gms/ltr.		1938
1F009	Fungal Agar w/low pH (Mycological Agar w/low pH) for selective enumeration and cultivation of saprophytic fungi and aciduric bacteria.	35.00 gms/ltr.		1938
1F010	Fungal Broth (Mycological Broth) for cultivation of fungi.	50.00 gms/ltr.		1827
1F011	*Fungobiotic Agar (Mycobio Agar), Modified for isolation of dermatophytes and other pathogenic fungi.	35.55 gms/ltr.	3114	
1G008	G. C. Agar Base for the cultivation of neisseria gonorrhoeae. This medium is made selective by the addition of appropriate antibiotic mixture e.g. Thayer Martin or NYC.	38.50		2022
1G002	Gelatin Iron Agar for detecting gelatin liquefaction and hydrogen sulphide production.	159.00 gms/ltr.		1824
1G009	Gelatin Mannitol Salt Agar (Staphylococcus Agar No. 110) for selective isolation and differentiation of pathogenic <i>Staphylococci</i> .	149.50 gms/ltr.	426	1544
1G003	Gelatin Salt Agar cultivation and differentiation of <i>Vibrio</i> species from foods.	65.00 gms/ltr.		1992
1G004	Glucose Cysteine Agar Base w/ Thiamine with added blood or haemoglobin or hemin, it is used for cultivation and enumeration of <i>Pasteurella tularensis</i> .	58.00 gms/ltr.	459	1830
1G005	Glucose Salt Teepol Broth (Twin pack) recommended by BIS for enrichment of <i>Vibrio parahaemolyticus</i> and marine isolates. gms of part A+4ml of part B	48.00	415	1878
1G006	Glucose Yeast Peptone Agar for isolation of yeasts from soil specimens.	50.00 gms/ltr.		1803
1I003	Glycerol Asparagine Agar Base see ISP Medium No.5	23.00 gms/ltr.	1497	4557
1G001	GN Broth, Hajna for selective enrichment of gram-negative organisms of the enteric group.	39.00 gms/ltr.	351	1358
1H001	Hanahan's Broth (SOB Medium) for use in cultivation of recombinant strains of <i>Escherichia coli</i> .	28.00 gms/ltr.		1764
1H003	Heart Infusion Agar for isolation and cultivation of a wide variety of fastidious organisms.	40.00 gms/ltr.	424	1884
1H004	Heart Infusion Broth for isolation and cultivation of a wide variety of fastidious organisms.	25.00 gms/ltr.	424	1884
1H002	Heart Infusion Powder, a rich nutritive component used in media employed for cultivation of fastidious organisms and antibiotic sensitivity.			2080

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. MEDIUM	100 GM	500 GM
1H005	Hektoen Enteric Agar for differential and selective isolation of <i>Salmonella</i> and <i>Shigella</i> species from enteric pathological specimens.	76.67 gms/ltr.	361	1605
1H006	Hemorrhagic Coli (HC) Agar for isolation & enumeration with an enzyme labeled monoclonal antibody of <i>Escherichia coli</i> .	61.13 gms/ltr.		1938
1H007	Hoyle Medium Base a highly selective medium used for the isolation and differentiation of <i>Corynebacterium diphtheriae</i> types.	40.00 gms/ltr.	424	1884
1H008	Hugh Leifson Glucose Medium for differentiation of <i>Staphylococci</i> from <i>Micrococci</i> on the basis of anaerobic fermentation of glucose.	45.52 gms/ltr.	424	1926
1H009	Hugh Leifson Medium recommended by BIS for detecting aerobic and anaerobic breakdown of glucose.	20.33 gms/ltr.	441	1994
1I004	Indole Nitrate Medium (Tryptone Nitrate Medium) for identification of microorganisms by means of nitrate reduction and indole production.	25.00 gms/ltr.	343	1416
1I009	Inhibitory Mould Agar, Ulrich (Mould Inhibitory Agar, Ulrich) for selective isolation of pathogenic fungi.	36.17 gms/ltr.		1884
1I005	Inorganic Salt Medium studying soil microorganisms such as <i>Rhizobium</i> species.	4.73 gms/ltr.	406	1827
1I006	Inorganic Salt Starch Agar see ISP Medium No 4	37.00 gms/ltr.	1108	3537
1I007	Inositol Gelatin Medium for the cultivation of <i>Plesiomonas shigelloides</i> from foods in accordance with APHA.	140.00 gms/ltr.		1917
1I008	Isolation Medium For Iron Bacteria for the isolation of iron bacteria, especially those belonging to <i>Sphaerotilus-Leptothrix</i> group.	10.90 gms/ltr.	406	1830
1I001	ISP Medium No.1 (Tryptone Yeast Extract Broth) a general purpose medium for not particularly fastidious microorganisms.	8.00 gms/ltr.	1045	3336
1I002	ISP Medium No.2 (Yeast Malt Agar) for the isolation and cultivation of yeasts, moulds and other aciduric microorganisms.	41.00 gms/ltr.	993	2997
1I006	ISP Medium No 4 (Inorganic Salt Starch Agar) for cultivation of <i>Streptomyces</i> species in accordance with International Streptomyces Project	37.00 gms/ltr.	1108	3537
1I003	ISP Medium No.5 (Glycerol Asparagine Agar Base) for cultivation of <i>Streptomyces</i> species as per International Streptomyces Project.	23.00 gms/ltr.	1497	4557
1K001	KF Streptococcai Agar Base for selective isolation and enumeration of faecal <i>Streptococci</i> in surface water by direct plating or by membrane filter method.	76.40 gms/ltr.		1724
1K002	KF Streptococcai Broth for detection and enumeration of faecal <i>Streptococci</i> in waters and examination of faces & other materials.	57.00 gms/ltr.		1626
1K003	KF Streptococcus Agar w/ BCP for detection and enumeration of faecal <i>Streptococci</i> .	76.40 gms/ltr.	572/5 vi	2080
1K004	KF Streptococcus Broth w/ BCP for detection and enumeration of faecal <i>Streptococci</i> .	56.40 gms/ltr.	572/5 vi	1962

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. MEDIUM	100 GM	500 GM
1K005	King's of Medium Base for studying oxidation-fermentation of carbohydrates by <i>Campylobacter species</i> .	0.50 gms/ltr.	366	1648
1K006	Kligler Iron Agar recommended by ISO Committee for identification of <i>Pseudomonas</i> species. It can be also used for differential identification of gram-negative enteric bacilli on the Basis of the fermentation of dextrose, lactose and H ₂ O production.	57.70 gms/ltr.		1637
1K007	Koser Citrate Medium to differentiate <i>Escherichia coli</i> & <i>Enterobacter aerogenes</i> on the basis of citrate utilization.	5.70 gms/ltr.	369	1515
1B003	L. Leichmannii Maintenance Medium (See *B12 Culture Agar)	42.10 gms/ltr.	653	
1E016	Lactobacilli Broth (See Elliker Broth)	48.50 gms/ltr.		1707
1L001	Lactose Broth recommended by BIS for the detection of coliform bacteria in water, foods, dairy products. in accordance with Standard Methods.	13.00 gms/ltr.	243	1152
1L002	Lactose Gelatin Medium for the detection of Clostridium species from food samples.	135.00 gms/ltr.		1828
1L003	Lauryl Sulphate Broth (Lauryl Tryptose Broth) for detection and enumeration of coliform bacteria in water, wastewater, dairy products & other foods.	35.60 gms/ltr.	309	1350
1L004	Lauryl Tryptose Mannitol Broth w/ Tryptophan a single tube medium used for confirmation of <i>Escherichia coli</i> in drinking water.	35.80 gms/ltr.		1860
1L005	Legionella Enrichment Broth Base with addition of supplements it is used for cultivation of <i>Legionella</i> species.	27.50 gms/ltr.	2115	
1L006	*Letheen Agar to determine the phenol coefficient of quaternary ammonium compounds using <i>Escherichia coli</i> or <i>Staphylococcus aureus</i> ATCC 6538.	32.00 gms/ltr.		2196
1L017	*Letheen Agar, Modified (*Modified Letheen Agar) for screening cosmetic products for microbial contamination.x	54.00 gms/ltr.		2196
1L007	*Letheen Broth, Modified for screening cosmetic products for microbial contamination in accordance with FDA.	42.80 gms/ltr.		2196
1L008	*Listeria Enrichment Broth, Modified for selective enrichment of <i>Listeria</i> species.	52.00 gms/ltr.	687	3128
1L009	*Listeria Enrichment Medium for selective isolation and cultivation of <i>Listeria Mono cutogenes</i> from clinical specimens.	54.37 gms/ltr.	687	3128
1L010	Listeria Motility Medium recommended by ISO committee for testing motility of <i>Listeria monocytogenes</i> .	29.60 gms/ltr.		1828
1L011	Lowenstein Jensen Medium Base (L.J. Medium) for isolation and cultivation of <i>Mycobacterium</i> species.	37.24 gms/ltr.	348	1464
1L012	Lysine Arginine Iron (LAL) Agar for the isolation and presumptive identification of <i>Yersinia</i> species from milk and milk products.	44.56 gms/ltr.	1464	
1L013	Lysine Decarboxylase Broth for differentiating <i>Salmonella arizonae</i> from the Bethesda Bellerup group of <i>Enterobacteriaceae</i> .	14.00 gms/ltr.	401	1513

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1L014	Lysine Decarboxylase Broth w/o Peptone recommended by ISO Committee for distinguishing the Salmonella arizonae from the Bethesda Ballerup group of Enterobacteriaceae.	9.00 gms/ltr.	390	1513
1L015	Lysine Iron Agar for differentiation of enteric organisms especially Salmonella and Arizona species, based on their ability to decarboxylate or deaminate lysine & to form hydrogen sulphide (H ₂ S).	34.56 gms/ltr.	370	1485
1L016	Lysine Medium Base isolation and enumeration of wild yeasts in pitching yeasts.	66.00 gms/ltr.	424	1780
1M009	MacConkey Agar recommended by BIS for isolation and differentiation of lactose fermenting & lactose nonfermenting enteric bacteria.	55.07 gms/ltr.	297	1443
1M005	MacConkey Agar w/ 0.15% Bile Salts, CV and NaCl for selective isolation and differentiation of coliform organisms & other enteric pathogens.	51.50 gms/ltr.	280	1234
1M006	MacConkey Agar w/CV, NaCl and 0.15% Bile Salts to identify Enterobacteriaceae in the presence of coliforms & lactose nonfermenters from water, sewage, food products etc.	51.55 gms/ltr.		1287
1M007	MacConkey Agar w/ 1.35% Agar, CV, NaCl and 0.15% Bile Salts for selective isolation & differentiation of lactose fermenting & lactose nonfermenting enteric bacteria, in accordance with B.P./U.S.P.	50.00 gms/ltr.	280	1287
1M044	MacConkey Agar w/ Bromo Thymol Blue for selective differentiation of lactose fermenting and non- fermenting enteric bacteria.	51.53 gms/ltr.	384	1513
1M008	MacConkey Agar w/o CV w/ 0.15% Bile Salts for selective isolation and differentiation of lactose fermenting and lactose nonfermenting enteric bacteria.	51.53 gms/ltr.	280	1234
1M010	MacConkey Broth w/ Neutral Red recommended by BIS for selective enrichment and enumeration of coliforms.	40.07 gms/ltr.	288	1234
1M011	MacConkey Broth Purple w/BCP for the presumptive identification of coliforms from variety of specimens such as water, milk and food etc.	40.00 gms/ltr.	280	1152
1S017	MacConkey Sorbitol Agar (see Sorbitol Agar)	50.00 gms/ltr.		2205
1M014	Malonate Broth, Ewing Modified for differentiation of members of Enterobacteriaceae on the basis of malonate utilization.	9.30 gms/ltr.	2070	
1M015	Malt Agar for detection and isolation of yeasts and moulds from dairy products, foods and other materials. Also for carrying stock cultures of yeasts and moulds.	45.00 gms/ltr.		2337
1M016	Malt Extract Agar Base (Malt Extract Agar w/Mycological Peptone) for detection, isolation and enumeration of yeasts & moulds.	50.00 gms/ltr.		2224
1M017	Malt Extract Agar Base, Modified as per Thorn and Church for isolation and cultivation of yeasts and moulds.	31.28 gms/ltr.		2337
1M018	Malt Extract Broth for detection and enumeration of yeasts, moulds and aciduric microorganisms.	20.00 gms/ltr.		2205
1M045	Maltodextrin Powder			850
1M013	Malonate Broth for differentiation of <i>Enterobacter</i> and <i>Escherichia</i> on the basis of malonate utilization.	8.00 gms/ltr.	1917	

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1M019	Mannitol Salt Agar for selective isolation of pathogenic <i>Staphylococci</i> .	111.00 gms/ltr.	325	1396
1M020	Mannitol Salt Broth for selective isolation of presumptive pathogenic <i>Staphylococci</i> .	96.00 gms/ltr.		1828
1M021	Middlebrook 7H9 Agar Base A for isolation, cultivation and sensitivity testing of <i>Mycobacterium tuberculosis</i> .	19.60 gms/ltr.		1938
1M022	Middlebrook 7H9 Broth Base for cultivation and sensitivity testing of <i>Mycobacterium tuberculosis</i> .	4.70 gms/ltr.		1828
1M023	Middlebrook 7H10 Agar Base for isolation, cultivation and sensitivity testing of <i>Mycobacterium tuberculosis</i> .	19.47 gms/ltr.		1938
1M024	Middlebrook 7H11 Agar Base for isolation, cultivation and sensitivity testing of <i>Mycobacteria</i> .	20..50 gms/ltr.		1938
1M025	Middlebrook 7H11 Agar Base w/o Malachite Green for isolation, cultivation and determination of antimicrobial susceptibility of <i>Mycobacteria</i> .	20.50 gms/ltr.		1938
1M001	MIO Medium (Motility Indole Ornithine Medium) for the identification of <i>Enterobacteriaceae</i> on the basis of motility, indole production & ornithine decarboxylase activity.	31.00 gms/ltr.		1824
1M002	MIU Medium Base for detection of motility, urease and indole production.	18.00 gms/ltr.		2034
1M026	Modified Duncan Strong (DS) Medium isolation & differentiation of <i>Clostridium perfringens</i> from other <i>Clostridia</i> from foods on the basis of raffinose fermentation.	34.00 gms/ltr.		3461
1L017	*Modified Lethen Agar (see Lethen Agar, Modified)	54.00 gms/ltr.		2196
1L007	*Modified Lethen Broth (see Lethen Broth, Modified)	42.80 gms/ltr.		2070
1M029	Modified Rappaport Vassiliadis Medium for selective enrichment of <i>Salmonellae</i> from food & environmental specimens.	30.00 gms/1110ml		1860
1M030	*Modified Rogosa Agar (M16 Agar) for cultivation, enumeration of lactic <i>Streptococci</i> used in manufacturing of cheddar cheese.	36.00 gms/ltr.		1823
1M032	Motility Medium S Base for easy detection of bacterial motility by means of TTC reduction.	60.00 gms/ltr.		1617
1M033	Motility Nitrate Medium, Buffered for isolation and detection of <i>Clostridium perfringens</i> on the basis of motility and nitrate test.	23.50 gms/ltr.		1938
1M034	Motility Nitrate Medium, Buffered recommended by ISO Committee for isolation & detection of <i>Clostridium perfringens</i> on the basis of motility & nitrate test.	19.50 gms/ltr.		1938
1M035	Motility Test Medium for detection of bacterial motility.	20.00 gms/ltr.		1688
1M036	Motility Test Medium (Edwards and Ewing) for testing motility of enteric bacteria.	22.00 gms/ltr.		1823
1I009	Mould Inhibitory Agar, Ulrich (see Inhibitory Mould Agar, Ulrich)	36.17 gms/ltr.		1884

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1M003	MR-VP Medium for performance of Methyl Red & Voges Proskauer tests in differentiation of coliaerogenesgroup.	17.00 gms/ltr.	226	909
1M004	MR-VP Medium recommended by BIS for performance of Methyl Red and Voges Proskauer tests in differentiation of coli-aerogenes group.	15.00 gms/ltr.	249	1000
1M037	Mueller Hinton Agar for cultivation of <i>Neisseria</i> and for determination of susceptibifity of microorganisms to antimicrobial agents.	38.00 gms/ltr.	333	1490
1M038	Mueller Hinton Broth to determine the susceptibility of bacteria to <i>Sulphonamides</i> by the tube dilution method.	21.00 gms/ltr.	401	1653
1F011	Mycobio Agar, Modified (See *Fungobiotic Agar)	35.55 gms/ltr.	3114	
1F008	Mycological Agar see Fungal Agar	35.00 gms/ltr.		1930
1M040	Mycological Agar, Modified for cultivation of fungi.	36.00 gms/ltr.		2024
1F009	Mycological Agar w/low pH (see Fungal Agar w/low pH)	35.00 gms/ltr.		1938
1F010	Mycological Broth see Fungal Broth	50.00 gms/ltr.		1890
1P038	Mycological Peptone (see Peptone M)			1548
1M043	Mycoplasma Agar Base (PPLO Agar Base) with the addition of enrichment it is used for isolation and cultivation of Mycoplasmas species (Pleuro-pneumonia-like organisms-PPLO)	36.00 gms/ltr.	484	1938
1A026	Neomycin, Erythromycin Assay Agar (Erythromycin Seed Agar) (see Antibiotic Assay Medium No. 11)	30.50 gms/ltr.	442	1769
1B017	Nickerson Medium (see Bi. G.G.Y. Agar)	45.00 gms/ltr.		1917
1A009	NIH Thioglycollate Broth (See Alternative Thioglycollate Medium)	29.00 gms/ltr.	183	699
1N003	Nitrate Agar for detection of nitrate reducing bacteria.	21.00 gms/ltr.		1938
1N004	Nitrate Broth for detection of nitrate reduction by bacteria. Also recommended by ISO Committee for the enumeration of <i>Bacillus cereus</i> .	9.00 gms/ltr.	406	1828
1N005	Nutrient Agar for cultivation of less fastidious microorganisms, can be enriched with blood or other biological fluids.	28.00 gms/ltr.	280	1242
1N008	Nutrient Agar, pH 6.0 w/0.8% NaCl for cultivation of bacteria requiring slightly acidic pH.	31.00 gms/ltr.	345	1464
1N006	Nutrient Agar, pH 6.8 a general purpose nutrient medium for examination of water,sewage, faeces and other materials.	23.00 gms/ltr.	345	1464

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1N007	Nutrient Agar, pH 7.0 recommended by ISO Committee for cultivation of <i>Salmonella</i> species.	23.00 gms/ltr.		1566
1N009	Nutrient Broth for general cultivation of less fastidious microorganisms, can be enriched with blood or other biological fluids	13.00 gms/ltr.	262	1081
1N010	Nutrient Gelatin for detection of gelatin liquefaction by proteolytic microorganisms.	128.00 gms/ltr.		1486
1O001	Ornithine Decarboxylase Broth recommended by ISO Committee for detection of the ability of microorganisms to decarboxylate ornithine.	9.00 gms/ltr.	1356	
1P039	Pancreatic Digest of Casein			1700
1P037	Pancreatic Digest of Gelatin			1585
1P037	Peptic Digest of Soyabean Meal			1918
1P034	Peptone, Bacteriological contains high tryptophan content used as culture media ingredient in variety of media. Also useful for commercial production of enzymes, vaccines, antibiotics and other products.			1161
1P038	Peptone M			1548
1P001	Peptone Sorbitol Bile Broth for identification of <i>Yersinia enterocolitica</i> from dairy products.	30.93 gms/ltr.		1884
1P035	Peptone Water as a growth medium and as the basis of carbohydrate fermentation media.	15.00 gms/ltr.	306	1229
1P008	Phenol Red Broth Base a basal medium to which carbohydrates are added for determination of fermentation reactions of pure cultures of microorganisms.	16.00 gms/ltr.		1605
1P009	Phenol Red Broth Base w/ Meat Extract highly nutritive basal medium which can be used to study fermentation of carbohydrates.	16.00 gms/ltr.		1605
1P002	Phenol Red Dextrose Agar for dextrose fermentation studies of microorganisms.	41.00 gms/ltr.		1702
1P010	Phenol Red Dextrose Broth for dextrose fermentation studies of microorganisms.	21.00 gms/ltr.		1605
1P011	Phenol Red Dulcitol Broth for dulcitol fermentation studies of microorganisms.	21.00 gms/ltr.	3186	
1P003	Phenol Red Lactose Agar for lactose fermentation studies of microorganisms.	41.00 gms/ltr.		1702
1P012	Phenol Red Lactose Broth recommended by ISO committee for lactose fermentation studies of coliforms.	25.00 gms/ltr.		1605
1P004	Phenol Red Maltose Agar for maltose fermentation studies of microorganisms.	41.00 gms/ltr.		1938
1P013	Phenol Red Maltose Broth for maltose fermentation studies of microorganisms.	21.00 gms/ltr.		1828
1P005	Phenol Red Mannitol Agar for mannitol fermentation studies of microorganisms.	41.00 gms/ltr.		1938

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1P014	Phenol Red Raffinose Broth for raffinose fermentation studies of microorganisms.	21.00 gms/ltr.	1828	
1P015	Phenol Red Rhamnose Broth for rhamnose fermentation studies of microorganisms.	21.00 gms/ltr.	3522	
1P016	Phenol Red Salicin Broth for salicin fermentation studies of microorganisms.	21.00 gms/ltr.	2841	
1P017	Phenol Red Sorbitol Broth for sorbitol fermentation studies of microorganisms.	21.00 gms/ltr.	987	
1P018	Phenol Red Starch Broth for starch fermentation studies of microorganisms.	21.00 gms/ltr.	860	
1P006	Phenol Red Sucrose Agar for sucrose fermentation studies of microorganisms.	41.00 gms/ltr.		1702
1P019	Phenol Red Sucrose Broth for sucrose fermentation studies of microorganisms.	21.00 gms/ltr.		1605
1P007	Phenol Red Tartrate Agar for identification and differentiation of <i>salmonellae</i> on the basis of tartrate utilization.	40.00 gms/ltr.	1377	
1P020	Phenol Red Trehalose Broth for trehalose fermentation studies of microorganisms.	21.00 gms/ltr.	3522	
1P022	Plate Count Agar for determining plate counts of microorganisms in milk & dairy products by pour plate technique.	17.50 gms/ltr.	405	1704
1P023	Plate Count Agar recommended by BIS for determining plate counts of microorganisms in milk and dairy products by pour plate technique.	30.00 gms/ltr.	401	1823
1P024	Plate Count Agar, Special for estimation of microbial counts in raw milk and other dairy products in accordance with Netherlands Dairy Association.	40.50 gms/ltr.		2931
1P021	Plate Count Agar (Standard Methods Agar) for determination of plate counts of microorganisms in foods, water and wastewater.	23.50 gms/ltr.	361	1517
1A025	Polymyxin Seed Agar (See Antibiotic Assay Medium No. 10)	52.00 gms/ltr.		2019
1P025	Potato Dextrose Agar for isolation and enumeration of yeasts and moulds from dairy and other food products.	39.00 gms/ltr.	321	1431
1P026	Proteose Peptone, Certified a highly nutritious ingredient employed in media used for bulk production of antibiotics, enzymes, bacterial toxins etc.			3321
1P027	Protose enzymic digest of mixed proteins, recommended for fermentation and vaccine industries.		1590	
1P029	Pseudomonas Agar (For Fluorescein) for detection of fluorescein production by <i>Pseudomonas</i> species.	38.00 gms/ltr.	387	1503
1P030	Pseudomonas Agar (For Fluorescein) for detection of fluorescein production by <i>Pseudomonas</i> species in accordance with U.S.P.	38.00 gms/ltr.	435	1747
1P031	Pseudomonas Agar (For Pyocyanin) for detection of pyocyanin production by <i>Pseudomonas</i> species.	46.40 gms/ltr.	388	1503

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1P032	Pseudomonas Agar (For Pyocyanin) for detection of pyocyanin production by <i>Pseudomonas</i> species in accordance with I.P./U.S.P.	46.40 gms/ltr.	435	1747
1P036	Pseudomonas Asparagine Broth for presumptive determination of <i>Pseudomonas aeruginosa</i> from recreational or natural water in accordance with A.P.H.A.	4.50 gms/ltr.	1593	5562
1P028	Pseudomonas Isolation Agar for selective isolation and identification of <i>Pseudomonas aeruginosa</i> from clinical and nonclinical specimens.	45.03 gms/ltr.	392	1752
1P033	Purple Broth Base for preparation of carbohydrate media used in fermentation studies for the cultural identification of pure cultures of enteric and other microorganisms.	15.00 gms/ltr.		1905
1R001	Rappaport Vassiliadis Medium for enrichment of <i>Salmonellae</i> , based on its ability to multiply selectively at high osmotic pressure, low pH and at 43° C, with modest nutritional requirements.	49.20 gms/ltr.	423	1905
1R002	Reinforced Clostridial Agar for the cultivation and enumeration of <i>Clostridia</i> and other anaerobes.	51.00 gms/ltr.	424	1695
1R003	Reinforced Clostridial Broth for the cultivation & enumeration of <i>Clostridia</i> & other anaerobes in accordance with B.P /E.P	38.00 gms/ltr.s	424	1695
1C022	Robinson's Cooked Meat Medium (R.C. Medium) see Cooked Meat Medium	125.00 gms/ltr.	433	1770
1R005	*Rogosa SL Agar for selective cultivation of oral and faecal <i>Lactobacilli</i> .	75.00 gms/ltr.		1863
1R006	*Rogosa SL Broth for selective cultivation of all <i>Lactobacilli</i> including oral & faecal <i>Lactobacilli</i> .	60.00 gms/ltr.		1756
1S007	Sabouraud Dextrose Agar for cultivation of yeasts, moulds and aciduric microorganisms in accordance with U.S.P.	65.00 gms/ltr.	345	1464
1D012	Sabouraud Dextrose Agar Base, Modified (See Dextrose Agar Base, Emmons)	47.00 gms/ltr.	451	1828
1S008	Sabouraud Maltose Agar for propagation of yeasts and moulds, particularly the parasitic fungi concerned with skin and scalp lesions.	65.00 gms/ltr.		1764
1S009	Sabouraud Maltose Broth for propagation of yeasts and moulds, particularly the parasitic fungi concerned with skin and scalp lesions.	50.00 gms/ltr.		1764
1F003	Sabouraud Medium, Fluid (see Fluid Sabouraud Medium)	30.00 gms/ltr.	190	748
1S011	Salmonella Agar, ONOZ for selective isolation & identification of <i>Salmonellae</i> from clinical specimens.	80.31 gms/ltr.		1887
1A020	Seed Agar (See Antibiotic Assay Medium No.1)	30.50 gms/ltr.	442	1769
1S035	Selenite Broth recommended as asn enrichment media for the isolation of <i>Salmonellae</i> .	23.00 gms/ltr.	498	1999
1S013	Selenite Broth Base w/o Selenite with addition of selenite, it is used for enrichment of <i>Salmonellae</i> from food, dairy products and pathological materials.	19.00 gms/ltr.		1701

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. MEDIUM	100 GM	500 GM
1F004	Selenite Cystine Broth (see Fluid Selenite Cystine Medium)	23.00 gms/ltr.	498	1999
1S012	Selenite F Broth an enrichment medium for isolation of <i>Salmonella</i> from faeces, urine or other pathological materials, in accordance with I.P.	23.00 gms/ltr.	484	1940
1S001	SF Broth for selective cultivation, detection and differentiation of Enterococci from other cocci in diagnostic work.	36.00 gms/ltr.		1560
1S002	S.F.P. Agar Base for the presumptive identification and enumeration of <i>Clostridium perfringens</i> in foods.	47.00 gms/ltr.		2058
1S003	SIM Medium for determination of hydrogen sulphide production, indole formation and motility of enteric bacilli.	36.23 gms/ltr.		1473
1S015	Simmons Citrate Agar for differentiating members of Enterobacteriaceae on the basis of citrate utilization.	24.28 gms/ltr.	363	1518
1S016	Sodium Tauroglycocholate		840	3224
1S017	Sorbitol Agar (Sorbitol MacConkey Agar) for isolation and identification of enteropathogenic <i>Escherichia coli</i> strains associated with infant diarrhoea.	50.00 gms/ltr.		2205
1S018	Sorbitol Iron Agar for cultural identification and differentiation of enteropathogenic <i>Escherichia coli</i> which do not ferment sorbitol.	46.00 gms/ltr.		2034
1S034	Soya Lecethin Paste		410	1517
1S019	Soyabean Casein Digest Agar (Tryptone Soya Agar) (Antibiotic Assay Medium NO. 36) a general purpose medium used for cultivation of a wide variety of microorganisms.	40.00 gms/ltr.	384	1539
1S020	Soyabean Casein Digest Medium (Tryptone Soya Broth) (Antibiotic Assay Medium NO. 37) a general purpose medium used for cultivation of a wide variety of microorganisms and sterility testing of moulds & lower bacteria.	30.00 gms/ltr.	181	699
1S021	Soyabean Casein Digest Medium a general purpose medium used for cultivation of a wide variety of microorganisms and sterility testing of moulds & lower bacteria in accordance with I.P.	30.00 gms/ltr.	181	861
1S010	Soyabean Casein Digest Medium (Tryptone Soya Broth w/o Dextrose) recommended for cultivation of anaerobic microorganisms when the presence of carbohydrate is not desired.	27.50 gms/ltr.		1458
1S014	Soyabean Casein Digest Medium w/ 0.1% Agar (Tryptone Soya Broth w/0.1% Agar) for cultivation of anaerobes from root canals, blood and other specimens.	31.00 gms/ltr.		1356
1A030	Sporulating Agar (Arret & Kirshbaum Medium) (See AK Agar No. 2)	30.80 gms/ltr.		1936
1S004	*SPS Agar for detection of <i>Clostridium perfringens</i> in foods.	40.00 gms/ltr.		1677
1S005	*SPS Agar Modified for selective isolation and enumeration of <i>Clostridium perfringens</i> from foodstuffs.	41.28 gms/ltr.		1677

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1S006	SS Agar (Salmonella Shigella Agar) for differential and selective isolation of <i>Salmonella</i> and <i>Shigella</i> species from pathological specimens, suspected foodstuffs etc.	63.00 gms/ltr.	345	1464
1P021	Standard Methods Agar (See Plate Count Agar)	23.50 gms/ltr.	361	1524
1S022	*Standard Methods Agar w/Tween 80 and Lecithin for sanitary examination of surfaces, that is, for counts before and after application of disinfectants.	29.20 gms/ltr.	1210	
1S023	Standard Nutrient Agar No.1 for the cultivation of fastidious bacteria.	37.00 gms/ltr.		1767
1S024	Standard Nutrient Broth No.1 for the cultivation of fastidious bacteria.	25.00 gms/ltr.		1668
1G009	Staphylococcus Agar No. 110 (see Gelatin Mannitol Salt Agar)	149.50 gms/ltr.	424	1545
1S026	Staphylococcus Agar No. 110 w/Azide for selective isolation and testing of pathogenic <i>Staphylococci</i> .	149.60 gms/ltr.		1926
1S027	Starch Agar for detection of starch hydrolyzing microorganisms.	30.00 gms/ltr.		1767
1A035	Stock Culture Agar (see Ayers and Johnson Agar)	50.00 gms/ltr.		1872
1A024	Streptomycin Assay Agar w/Yeast Extract (see Antibiotic Assay Medium No.5)	25.50 gms/ltr.	525	2018
1S030	*Sulphate API Agar w/o Sodium Lactate for detection and estimation of sulphate reducing bacteria.	25.40 gms/ltr.		2017
1S031	*Sulphate API Broth w/o Sodium Lactate for detection of sulphate reducing bacteria.	11.40 gms/ltr.		1905
1S032	*Sulphate API Broth w/o NaCl for detection, differentiation and estimation of sulphate reducing bacteria.	1.40 gms/ltr.	424	1905
1T001	TCBS Agar (Selective) for selective isolation of <i>Vibrio cholerae</i> and other enteropathogenic <i>Vibrios</i> .	89.00 gms/ltr.	441	1828
1T002	Tellurite Blood Agar Base for the selective isolation & cultivation of <i>Corynebacterium</i> species.	31.00 gms/ltr.		1918
1T003	Tetrathionate Brilliant Green Bile Broth for isolation and identification of <i>Salmonellae</i> in accordance with I. P.	63.00 gms/ltr.		1140
1T004	Tetrathionate Broth Base an enrichment broth for isolation of <i>Salmonellae</i> from specimens to be contaminated with <i>Salmonellae</i> in accordance with I. P.	77.40 gms/ltr.	285	1211
1F006	Tetrathionate Broth Base w/o Iodine and BG (See Fluid Tetrathionate Medium w/o Iodine and BG)	46.00 gms/ltr.	248	1080
1T005	Thayer Martin Medium Base for selective isolation of <i>Gonococci</i> from pathological specimens.	42.00	384	1605
1T006	Thiobacillus Agar for isolation and cultivation of <i>Thiobacillus</i> species.	22.70 gms/ltr.		1938
1T007	Thiobacillus Broth for cultivation of <i>Thiobacillus</i> .	10.16 gms/ltr.		1828

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. O MEDIUM	100 GM	500 GM
1A009	Thioglycollate Broth, Alternative (see Alternative Thioglycollate Medium)	29.00 gms/ltr.	181	699
1F007	Thioglycollate Medium Fluid (see Fluid Thioglycollate Medium)	29.75 gms/ltr.	181	699
1T010	Thioglycollate Medium w/o Dextrose for cultivation of aerobes, microaerophiles, anaerobes and for fermentation studies with various carbohydrates.	25.70 gms/ltr.		1269
1D024	Thioglycollate Medium w/o Indicator see Diagnostic Thioglycollate Medium	30.00 gms/ltr.		1356
1T035	Thiosulphate Citrate Broth a selective isolation medium for vibrio species particularly V. Cholera.	76.08 gms/ltr.		1873
1T012	Todd Hewitt Broth for cultivation of group A haemolytic <i>Streptococci</i> used for serological studies.	37.00 gms/ltr.	369	1607
1C004	Transport Medium w/o Charcoal see Cary-Blair Medium Base.	12.60 gms/ltr.	495	2093
1T014	Tributyryn Agar Base w/o Tributyrin for detection of lipolytic microorganisms.	23.00 gms/ltr.	424	1884
1T015	Triple Sugar Iron Agar for identification of gram-negative enteric bacilli on the basis of dextrose, lactose and sucrose fermentation and hydrogen sulphide production.	65.00 gms/ltr.	321	1365
1T040	Tryptone Agar for growth of phage.	33.00 gms/ltr.		1650
1T016	Tryptone Broth (Tryptone Water) for detection of indole producing microorganisms.	15.00 gms/ltr.	302	1290
1T017	Tryptone Glucose Beef Extract Agar (TGB Agar) for enumeration bacteria in water, air, milk & dairy products.	24.00 gms/ltr.	424	1918
1T018	Tryptone Glucose Extract Agar (Tryptone Glucose Yeast Extract Agar) for enumeration of bacteria in water, air, milk & dairy products.	24.00 gms/ltr.	327	1512
1T019	Tryptone Glucose Yeast Extract Broth for enumeration of microorganisms from foods by MPN technique.	17.25 gms/ltr.		1548
1I004	Tryptone Nitrate Medium (see Indole Nitrate Medium)	25.00 gms/ltr.	343	1416
1T021	Tryptone Phosphate Broth for enrichment and cultivation of enteropathogenic <i>Escherichia coli</i> from suspected food samples.	30.50 gms/ltr.		1683
1S019	Tryptone Soya Agar (Antibiotic Assay Medium No. 36) (see Soyabean Casein Digest Agar)	40.00 gms/ltr.	387	1539
1S020	Tryptone Soya Broth (Antibiotic Assay Medium No. 37) (see Soyabean Casein Digest Medium)	30.00 gms/ltr.	180	699
1S014	Tryptone Soya Broth w/0.1% Agar (see Soyabean Casein Digest Medium w/ 0.1% Agar)	31.00 gms/ltr.		1356
1S010	Tryptone Soya Broth w/o Dextrose (see Soyabean Casein Digest Medium)	27.50 gms/ltr.		1458

CODE	PRODUCT NAME	PRICE IN RUPEES		
		QTY. PER LTR. MEDIUM	100 GM	500 GM
1T026	Tryptone Tellurite Agar Base for selective isolation of pathogens from clinical specimens, especially from nose, throat & vagina.	47.00 gms/ltr.		1938
1T028	Tryptone Yeast Extract Agar recommended by ISO Committee for estimation of microbial counts in water.	21.00 gms/ltr.		2034
1I001	Tryptone Yeast Extract Broth (see ISP Medium No.1)	8.00 gms/ltr.	1045	3336
1T030	Tryptose Agar for isolation, cultivation and differentiation primarily of <i>Brucella</i> , but also of <i>Streptococci</i> , <i>Pneumococci</i> , <i>Meningococci</i> and other pathogenic microorganisms.	41.00 gms/ltr.		1938
1T031	Tryptose Blood Agar Base for the isolation fastidious organisms and determining the haemolytic reactions.	33.00 gms/ltr.	424	1803
1T032	Tryptose Broth for the cultivation primarily of <i>Brucella</i> species.	26.00 gms/ltr.		1526
1T033	Tryptose Phosphate Broth for cultivation of fastidious bacteria and as an adjuvant to tissue culture media.	29.50 gms/ltr.		1432
1U001	Urea Agar Base (Christensen) (Autoclavable) for detection of urease production, particularly by <i>Proteus vulgaris</i> , <i>Micrococci</i> and paracolon organisms.	24.00 gms/ltr.	378	1464
1U002	*Urea Broth (Filter Sterilizable) for identification of bacteria on the basis of urea utilization, specifically for the differentiation of <i>Proteus</i> species from <i>Salmonella</i> and <i>Shigella</i> species.	38.71 gms/ltr.		1626
1V001	Violet Red Bile Agar recommended by BIS for selective isolation, detection and enumeration of coli-aerogenes bacteria in water, milk, other dairy and food products.	41.53 gms/ltr.	357	1464
1V002	Violet Red Bile Agar (1.2%) recommended by ISO Committee for selective isolation & enumeration of coli-aerogenes in water, milk, other dairy & food products.	38.53 gms/ltr.		1464
1V003	Violet Red Bile Broth for detection and enumeration of coliforms from water and food.	26.53 gms/ltr.		1464
1B011	*Vitamin B12 Assay Medium (see *B12 Assay Medium)	85.00 gms/ltr.	1584	
1V005	Vogel-Johnson Agar Base w/o Tellurite (V.J. Agar) for selective isolation of coagulase positive, mannitol fermenting <i>Staphylococcus aureus</i> from heavily contaminated foods and clinical specimens.	61.00 gms/ltr.	363	1464
1W001	Wilson Blair Agar Base with the addition of selective reagent used for the isolation of <i>Salmonella typhi</i> .	60.00 gms/ltr.		1938
1W002	Wilson Blair Agar w/BG for isolation and preliminary identification of <i>Salmonella typhi</i> from clinical specimens.	52.00 gms/ltr.		1938
1W003	Wort Agar for the cultivation and enumeration of yeasts.	48.30 gms/ltr.		1926
1W004	Wort Broth for cultivation and enumeration of yeasts.	33.28 gms/ltr.		1815
1X001	Xylose Lysine Agar Base for isolation and identification of pathogenic enteric bacilli.	45.00 gms/ltr.		1641

CODE	PRODUCT NAME	QTY.	PRICE IN RUPEES	
		PER LTR. MEDIUM	100 GM	500 GM
1X002	Xylose Lysine Deoxycholate Agar (XLD Agar) for selective isolation & enumeration of <i>Salmonella typhi</i> and other <i>Salmonella</i> species.	56.68 gms/ltr.	363	1464
1A023	Yeast Beef Agar (See Antibiotic Assay Medium No.4)	26.50 gms/ltr.		2039
1Y001	Yeast Extract Agar highly nutritive medium recommended for plate count of microorganisms in water.	23.00 gms/ltr.	297	1202
1I002	Yeast Malt Agar (YM Agar) see ISP Medium No.2	41.00 gms/ltr.	993	2997
1Y003	Yeast Malt Broth (YM Broth) for isolation and cultivation of yeasts, moulds and aciduric microorganisms.	21.00 gms/ltr.	401	1605
1Y004	Yeast Mannitol Agar w/1.5% Agar for cultivation, isolation and enumeration of soil microorganisms like <i>Rhizobium</i> species.	27.80 gms/ltr.		1803
1Y005	Yeast Mannitol Broth for cultivation of <i>Rhizobium</i> species.	12.80 gms/ltr.		1701

Please Contact us for custom synthesis & technical information / inquiries on any of Blulux Dehydrated Culture Media Products or for the availability of specific Blulux Dehydrated Culture Media not currently in the price list.

We offer

**Special
Prices**

for Exports & Bulk Requirements

©Gms/Ltr. figure indicates wt. of powder medium required to prepare 1 Ltr. of medium